arXiv:hep-ph/9905249v5 12 Jan 2001

BI-TP 99/12

Two-loop self-energy master integrals on shell

J. Fleischer [1], M. Yu. Kalmykov [2] and A. V. Kotikov [3]

Fakultät für Physik, Universität Bielefeld, D-33615 Bielefeld, Germany

Abstract

Analytic results for the complete set of two-loop self-energy master integrals on shell with one mass are calculated.

Keywords: Feynman diagram, Pole mass.

PACS number(s): 12.38.Bx

---

[1] E-mail: fleischer@physik.uni-bielefeld.de

[2] E-mail: misha@physik.uni-bielefeld.de
On leave of absence from BLTP, JINR, 1141980 Dubna (Moscow Region) Russia

[3] E-mail: kotikov@sunse.jinr.ru
Particle Physics Lab., JINR, 1141980 Dubna (Moscow Region) Russia

For a wide class of low-energy observables the dominant and the next-to-leading effects originate entirely from the gauge boson propagators. These corrections are sensitive to heavy virtual particles due to quadratic divergences of the corresponding diagrams and due to non-decoupling of the heavy particles within the Higgs mechanism for the generation of masses. Only recently have the two-loop sub-leading corrections to the $m_W - m_Z$ interdependence of order $O(G_F^2 m_t^2 m_Z^2)$ been calculated in an expansion in terms of the large top mass [1]. Moreover, the two-loop Higgs boson mass dependence associated with the fermion loop to Δr has also become available [2] [4]. These results allow to reduce the theoretical uncertainties originating from higher order effects of the basic physical quantities [4]. To improve the accuracy of the calculations, corrections of order $O(G_F^2 m_Z^4)$ must be taken into account, which require in particular the calculation of two-loop self-energy diagrams with only gauge bosons and massless particles. Keeping in mind this physical application, we present here analytical results for all two-loop on-shell self-energy master integrals with one mass. Another application of the given results is the calculation of the "naive" part of diagrams arising within the asymptotic expansion when the external momentum is on the mass shell of the particle with the large mass [5].

The aim of this letter is the analytical calculation of a full set of two-loop on-shell self-energy master integrals with one mass, Eqs.(1)-(6). The diagrams of this type are presented on Fig.1. It can be shown [6, 7], that the full set of master-integrals consists of the following ones: **F11111, F00111, F10101, F10110, F01100, F00101, F10100, F00001, V1111, V1001, J111** (see Fig.1) with all indices (powers of propagators) equal to 1, and two integrals of the **J011**-type: with indices 111 and 112, respectively (for the massless integrals see [8, 9]). The integration-by-parts method [9, 10] allows not only to reduce diagrams with higher indices to master-integrals, but also to calculate the master integrals themselves [5]. The main idea for the latter step is the following: a diagram with the index $(i+1)$ on a massive line can be represented as the derivative of the same diagram with index i on that line w.r.t. its mass squared. This representation is used in certain equations obtained by the integration by part method, yielding a differential equation (in general partial differential equations) for the initial diagram (with the index i) which can be solved analytically. This approach [13, 14] - to construct the (differential) relations between diagrams - is called the 'Differential Equation Method'. For some particular cases (where the number of the masses is small) these partial differential equations can be rewritten as ordinary differential equations [6].

To obtain the finite part of two-loop physical results one needs to know the **F**-type integrals up to the finite part, the **V**- and **J**-type integrals up to the ε-part, and the one-loop integrals up to the ε^2-part. In the last years various methods have been suggested and many analytical results for one-, two- and three-loop self-energy diagrams with different values of masses and external momenta have been obtained. In particular, the ε-part of one-loop integrals is given in Ref.[16], the finite parts of two-loop self-energy diagrams with different internal masses and arbitrary external momentum are given in Refs.[12],[14], [17]-[24], the ε-part of two-loop tadpole and sunset-type integrals at threshold (pseudothreshold) are calculated in Refs.[25]-[27]. Very efficient and elegant methods for numerical calculation of Feynman diagrams have been presented in Refs.[28],[29]. In many of these cases the authors have obtained one-dimensional integrals as well for arbitrary masses and momentum. We do not know, however, of attempts to reduce these integrals to generalized polylogarithms or other analytic expressions.

---

[4] Recently the accuracy of these two results has been tested in [3].

[5] One of the first calculations of this type for the Standard Model and QED was performed in Refs.[11, 12].

[6] An example where no reduction to an ordinary differential equation was made, may be found in [15].

We consider here the following master-integrals in Euclidean space-time with dimension $N = 4 - 2\varepsilon$:

$$\begin{aligned}
\mathbf{ONS}\{\mathcal{I}\mathcal{J}\}(i,j,m) &\equiv K^{-1}\int d^Nk P^{(i)}(k,\mathcal{I}m)\ P^{(j)}(k-p,\mathcal{J}m)\Big|_{p^2=-m^2},\\
\mathbf{J}\{\mathcal{I}\mathcal{J}\mathcal{K}\}(i,j,k,m) &\equiv K^{-2}\int d^Nk_1d^Nk_2P^{(i)}(k_1,\mathcal{I}m)P^{(j)}(k_1-k_2,\mathcal{J}m)\\
&\qquad\times\ P^{(k)}(k_2-p,\mathcal{K}m)\Big|_{p^2=-m^2},\\
\mathbf{V}\{\mathcal{I}\mathcal{J}\mathcal{K}\mathcal{L}\}(i,j,k,l,m) &\equiv K^{-2}\int d^Nk_1d^Nk_2P^{(i)}(k_2-p,\mathcal{I}m)\\
&\quad\times P^{(j)}(k_1-k_2,\mathcal{J}m)P^{(k)}(k_1,\mathcal{K}m)\ P^{(l)}(k_2,\mathcal{L}m)\Big|_{p^2=-m^2},\\
\mathbf{F}\{\mathcal{A}\mathcal{B}\mathcal{I}\mathcal{J}\mathcal{K}\}(a,b,i,j,k,m) &\equiv m^2K^{-2}\int d^Nk_1d^Nk_2P^{(a)}(k_1,\mathcal{A}m)\\
&\quad\times P^{(b)}(k_2,\mathcal{B}m)P^{(i)}(k_1-p,\mathcal{I}m)\\
&\quad\times P^{(j)}(k_2-p,\mathcal{J}m)\ P^{(k)}(k_1-k_2,\mathcal{K}m)\Big|_{p^2=-m^2},
\end{aligned}$$

where

$$K = \frac{\Gamma(1+\varepsilon)}{(4\pi)^{\frac{N}{2}}\,(m^2)^{\varepsilon}},\quad P^{(l)}(k,m) \equiv \frac{1}{(k^2+m^2)^l},$$

the normalization factor $1/(2\pi)^N$ for each loop is assumed, and $\mathcal{A},\mathcal{B},\mathcal{I},\mathcal{J},\mathcal{K} = 0,1$.

The finite part of most of the F-type master-integrals can be obtained from results of Ref.[24] in the limit $z \to 1$. Nevertheless, to take this limit is not an easy task in general. We have used this approach to obtain the results for **F10110** and **F00101**. The finite part of **F10101** is given in Refs.[17, 18], **F00001** and **F10100** are presented in Ref.[20]. These integrals we have verified by taking the above mentioned limit $z \to 1$ of results in Ref.[24]. **F11111** is calculated in Ref.[23] and we have checked it numerically. Instead of the usually taken **F01101** integral [17, 12] we consider **J111** as master integral. We recall the results of all master integrals for completeness. The last unknown master integral then remains to be **F00111**. This appears to be one of the most complicated integrals and below we present details of its calculation.

The finite part of the integrals of V- and J-type can be found in Refs.[20]-[21]. The calculation of the ε e (ε^2) parts of master integrals of this type have been performed by the differential equation method [13, 14].

The results for F-type master-integrals are follows:

$$\mathbf{F}\{\mathcal{A}\mathcal{B}\mathcal{I}\mathcal{J}\mathcal{K}\}(1,1,1,1,1,m) = a_1\zeta(3) + a_2\frac{\pi}{\sqrt{3}}S_2 + a_3 i\pi\zeta(2) + \mathcal{O}(\varepsilon), \tag{1}$$

and the coefficients $\{a_i\}$ are given in Table I:

TABLE I

	F11111	**F00111**	**F10101**	**F10110**	**F01100**	**F00101**	**F10100**	**F00001**
a_1	-1	0	-4	-1	0	-3	-2	-3
a_2	$\frac{9}{2}$	9	$\frac{27}{2}$	9	$\frac{27}{2}$	$\frac{27}{2}$	9	0
a_3	0	0	$\frac{1}{3}$	0	1	1	$\frac{2}{3}$	1

where [11, 17]

$$S_2 = \frac{4}{9\sqrt{3}}\mathrm{Cl}_2\left(\frac{\pi}{3}\right) = 0.260434137632\cdots.$$

Here we used the $m^2 - i\varepsilon$ prescription. The results for the remaining master integrals are the following ones:

$$\mathbf{V}\{\mathcal{IJKL}\}(1,1,1,1,m) = \frac{1}{2\varepsilon^2} + \frac{1}{\varepsilon}\left(\frac{5}{2} - \frac{\pi}{\sqrt{3}}\right) + \frac{19}{2} + \frac{b_1}{2}\zeta(2) - 4\frac{\pi}{\sqrt{3}} - \frac{63}{4}S_2 + \frac{\pi}{\sqrt{3}}\ln 3 + \varepsilon\left\{\frac{65}{2} + b_2\zeta(2) - b_3\zeta(3) - 12\frac{\pi}{\sqrt{3}} - 63S_2 + b_4\zeta(2)\ln 3 + \frac{9}{4}b_4 S_2\frac{\pi}{\sqrt{3}} + \frac{63}{4}S_2\ln 3 + 4\frac{\pi}{\sqrt{3}}\ln 3 - \frac{1}{2}\frac{\pi}{\sqrt{3}}\ln^2 3 - \frac{b_5}{2}\frac{\pi}{\sqrt{3}}\zeta(2) - \frac{21}{2}\frac{\mathrm{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}}\right\} + \mathcal{O}(\varepsilon^2), \qquad (2)$$

where the coefficients $\{b_i\}$ are listen in Table II:

TABLE II

	b_1	b_2	b_3	b_4	b_5
V1111	-1	-6	$\frac{9}{2}$	4	9
V1001	3	8	$-\frac{3}{2}$	0	21

$$\mathbf{J111}(1,1,1,m) = -m^2\left(\frac{3}{2\varepsilon^2} + \frac{17}{4\varepsilon} + \frac{59}{8} + \varepsilon\left\{\frac{65}{16} + 8\zeta(2)\right\} - \varepsilon^2\left\{\frac{1117}{32} - 52\zeta(2) + 48\zeta(2)\ln 2 - 28\zeta(3)\right\} + \mathcal{O}(\varepsilon^3)\right), \qquad (3)$$

$$\mathbf{J011}(1,1,2,m) = \frac{1-4\varepsilon}{2(1-2\varepsilon)(1-3\varepsilon)}\left(\frac{1}{\varepsilon^2} + 2\frac{\pi}{\sqrt{3}} - \frac{2}{3}\zeta(2) + \varepsilon\left\{8\frac{\pi}{\sqrt{3}} - \frac{2}{3}\zeta(2) - 6\frac{\pi}{\sqrt{3}}\ln 3 + \frac{2}{3}\zeta(3) + 27S_2\right\} + \mathcal{O}(\varepsilon^2)\right), \qquad (4)$$

$$\mathbf{J011}(1,1,1,m) = -\frac{m^2}{2}\frac{4-15\varepsilon}{(1-2\varepsilon)(1-3\varepsilon)(2-3\varepsilon)}\Bigg(\frac{1}{\varepsilon^2}+\frac{3}{2}\frac{\pi}{\sqrt{3}}$$
$$+\varepsilon\Bigg\{\frac{45}{8}\frac{\pi}{\sqrt{3}}-\frac{9}{2}\frac{\pi}{\sqrt{3}}\ln 3+\frac{81}{4}S_2\Bigg\}+\varepsilon^2\Bigg\{12-\zeta(2)-\frac{1863}{16}S_2-\frac{867}{32}\frac{\pi}{\sqrt{3}}$$
$$+\frac{207}{8}\frac{\pi}{\sqrt{3}}\ln 3+\frac{243}{4}S_2\ln 3-\frac{27}{4}\frac{\pi}{\sqrt{3}}\ln^2 3-21\frac{\pi}{\sqrt{3}}\zeta(2)-\frac{81}{2}\frac{\mathrm{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}}\Bigg\}$$
$$+\mathcal{O}(\varepsilon^3)\Bigg), \tag{5}$$

$$\mathbf{ONS11}(1,1,m) = \frac{1}{1-2\varepsilon}\left[\frac{1}{\varepsilon}-\frac{\pi}{\sqrt{3}}+\varepsilon\left\{\frac{\pi}{\sqrt{3}}\ln 3-9S_2\right\}\right.$$
$$\left.+\varepsilon^2\left\{9S_2\ln 3-\frac{1}{2}\frac{\pi}{\sqrt{3}}\ln^2 3-6\frac{\mathrm{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}}-3\frac{\pi}{\sqrt{3}}\zeta(2)\right\}+\mathcal{O}(\varepsilon^3)\right], \tag{6}$$

where [31]

$$\mathrm{Ls}_3\left(\frac{2\pi}{3}\right) = -2.14476721256949\cdots$$

(to our knowledge $\mathrm{Ls}_3\left(\frac{2\pi}{3}\right)$ has appeared for the first time in the calculation of the ε-part of two-loop tadpole integrals in Ref.[25] [7]). The expansion of J111(1,1,1,m) up to ε^4 terms is given in Ref.[30].

The above results were checked numerically. Padé approximants were calculated from the small momentum Taylor expansion of the diagrams [32]. The Taylor coefficients were obtained by means of the package [33] with the master integrals taken from [19]. In this manner high precision numerical results for the on-shell values of most of the diagrams were obtained [34]. Further we made use of the idea of Broadhurst [35] to apply the FORTRAN program **PSLQ** [36] to express the obtained numerical values in terms of a 'basis' of irrational numbers, which were predicted by our analytical calculation, i.e. the differential equation method.

Let us point out that the numbers we obtain are related to polylogarithms at the sixth root of unity and hence are in the same class of transcendentals obtained by Broadhurst [35] in his investigation of three-loop diagrams which correspond to a closure of the two-loop topologies considered here.

To demonstrate our method of calculation we present in what follows some details for the **F00111** integral. The basis of our approach is the differential equation method, i.e. we write a differential equation for **F00111**, the inhomogeneous term of which is expressed in terms of some 'simpler' integrals. Before turning to the final step, we discuss first of all these latter integrals. All calculations are performed off shell and only at the end we take the on-shell limit. The following extra notations are needed:

a) tadpoles:

$$\mathbf{T}_1(i,m) \;\; \equiv \;\; \int d^N k \; P^{(i)}(k,m) = \frac{\Gamma(i-\frac{N}{2})}{(4\pi)^{\frac{N}{2}}\Gamma(i)}\left(m^2\right)^{\frac{N}{2}-i},$$

---

[7] The definition is: $\mathrm{Ls}_3(x) = -\int_0^x \ln^2\left|2\sin\frac{\theta}{2}\right| d\theta$.

$$\mathbf{T}_2(i,j,k,m) \equiv \int d^Nk_1 d^Nk_2 P^{(i)}(k_1,m)P^{(j)}(k_2,m)P^{(k)}(k_1-k_2,m),$$

b) loops with one massive line:

$$\mathbf{L}(i,j,m) \equiv \int d^Nk\ P^{(i)}(k,0)P^{(j)}(k-p,m),$$

c) sunset diagrams with two massive lines:

$$\mathbf{J}(k,i,j,m) \equiv \int d^Nk_1 d^Nk_2 P^{(k)}(k_2-p,0)P^{(i)}(k_1,m)P^{(j)}(k_1-k_2,m), \tag{7}$$

d) two-loop diagrams with four propagators:

$$\begin{aligned}\mathbf{V_1}(i,j,k,l,m) &\equiv \int d^Nk_1 d^Nk_2 P^{(i)}(k_2-p,m)P^{(j)}(k_1-k_2,0)P^{(k)}(k_1,m)P^{(l)}(k_2,0),\\ \mathbf{V_2}(i,j,k,l,m) &\equiv \int d^Nk_1 d^Nk_2 P^{(i)}(k_2-p,0)P^{(j)}(k_1-k_2,m)P^{(k)}(k_1,m)P^{(l)}(k_2,m),\end{aligned}$$

e) two-loop diagrams with five propagators:

$$\begin{aligned}\mathbf{F}(a,b,i,j,k,m) &\equiv \int d^Nk_1 d^Nk_2 P^{(a)}(k_1,0)P^{(b)}(k_2,0)\\ &\quad\times P^{(i)}(k_1-p,m)P^{(j)}(k_2-p,m)P^{(k)}(k_1-k_2,m).\end{aligned}$$

1. *One-loop diagrams with one massive line.*
Because it is ultraviolet finite, the integral with shifted indices, $\mathbf{L}$(1,2,m), is more suitable for analytic calculations than the 'usual' master integral with all indices equal to one. The recurrence relations

$$\begin{aligned}\mathbf{L}(1,1,m)\cdot(N-3) &= -2m^2\mathbf{L}(1,2,m)-(p^2+m^2)\mathbf{L}(2,1,m),\\ \mathbf{L}(1,1,m)\cdot(N-3) &= \mathbf{T}_1(2,m)-(p^2+m^2)\mathbf{L}(1,2,m),\end{aligned} \tag{8}$$

allow to express the needed integrals in terms of the diagram $\mathbf{L}(1,2,m)$, which can be written as:

$$\begin{aligned}\mathbf{L}(1,2,m) &= \frac{K}{p^2}\int_0^1 ds\frac{1}{(1-s)^{\varepsilon}(1+as)^{(1+\varepsilon)}}\\ &= \frac{K}{p^2}\Bigg(\log(1+a)-\varepsilon\Big[\log^2(1+a)+\mathrm{Li}_2(-a)\Big]+\varepsilon^2\Big[\frac{2}{3}\log^3(1+a)\\ &\quad+2\log(1+a)\mathrm{Li}_2(-a)+2\mathrm{S}_{1,2}(-a)-\mathrm{Li}_3(-a)\Big]\Bigg)+O(\varepsilon^3),\end{aligned}$$

where $a=p^2/m^2$ (a = -1 corresponds to the on-shell case). To obtain this result we used the representation (see e.g. [24, 37])

$$\int d^N k P^{(i)}(k, m_1) P^{(j)}(k-p, m_2) = \frac{\Gamma(i+j-N/2)}{(4\pi)^{\frac{N}{2}}\Gamma(i)\Gamma(j)}$$
$$\times \int_0^1 ds \frac{1}{s^{(i+1-N/2)}(1-s)^{(j+1-N/2)}} P^{(i+j-N/2)}\left(p, \frac{m_1}{s} + \frac{m_2}{1-s}\right). \qquad (9)$$

2. *The sunset diagram with two massive lines.*
The set of master-integrals of this type consists of two integrals, e.g. **J011**(1,1,1,m) and **J011**(1,1,2,m). To avoid ultraviolet singularities, however, we choose again integrals with shifted indices: $\mathbf{J}(1,2,2,m)$ and $\mathbf{J}(1,1,3,m)$. Appropriate recurrence relations connecting these integrals need to be derived (see also Refs.[6, 38]). Applying the integration by parts relation with the massless line as distinctive one [8] , we get

$$\begin{aligned}
&\mathbf{J}(1,1,2,m)\cdot(N-2) \\
&= 2\int d^N k_1 d^N k_2 P^{(1)}(k_2-p,0)P^{(2)}(k_1,m)P^{(2)}(k_1-k_2,m)\cdot(k_2-k_1)^\mu(k_2-p)^\mu \\
&= \int d^N k_1 d^N k_2 P^{(1)}(k_2-p,0)P^{(2)}(k_1,m)P^{(2)}(k_1-k_2,m)\cdot k_2^\mu(k_2-p)^\mu, \qquad (10)
\end{aligned}$$

where for the last step the symmetry of the integral is taken into account. Expanding the scalar product $2(k_2, k_2-p) = k_2^2 + (p-k_2)^2 - p^2$, we have

$$\begin{aligned}
&\mathbf{J}(1,1,2,m)\cdot(N-2) = \frac{1}{2}\mathbf{T}_1^2(2,m) - \frac{p^2}{2}\mathbf{J}(1,2,2,m) \\
&+\frac{1}{2}\int d^N k_1 d^N k_2 P^{(1)}(k_2-p,0)P^{(2)}(k_1,m)P^{(2)}(k_1-k_2,m)\cdot k_2^2. \qquad (11)
\end{aligned}$$

Applying now the integration-by-parts relation with the massive line with index 2 as distinctive one, we obtain

$$\begin{aligned}
&\mathbf{J}(1,1,2,m)(N-6) = -4m^2\mathbf{J}(1,1,3,m) - 2m^2\mathbf{J}(1,2,2,m) \\
&-\int d^N k_1 d^N k_2 P^{(1)}(k_2-p,0)P^{(2)}(k_1,m)P^{(2)}(k_1-k_2,m)\cdot k_2^2. \qquad (12)
\end{aligned}$$

Combining Eq.(11) and Eq.(12) we get the final relation:

$$\mathbf{J}(1,1,2,m)2(1-3\varepsilon) = \mathbf{T}_1^2(2,m) - (p^2+2m^2)\mathbf{J}(1,2,2,m) - 4m^2\mathbf{J}(1,1,3,m). \qquad (13)$$

The second recurrence relation,

$$\begin{aligned}
\mathbf{J}(1,1,1,m)\,(1-2\varepsilon)\left(1-\frac{3}{2\varepsilon}\right) &= \left[\frac{\varepsilon}{2}p^2 - (2-5\varepsilon)\,m^2\right]\mathbf{J}(1,1,2,m) \\
&-m^4\mathbf{J}(1,2,2,m) - m^2(p^2+2m^2)\mathbf{J}(1,1,3,m), \qquad (14)
\end{aligned}$$

---

[8] In the integration-by-parts procedure one line is considered 'distinctive', i.e. its propagator contains nothing but the momentum k_μ which is used for the partial integration (see, e.g. the discussion in Ref.[24]).

can be obtained by applying the operator $\Big(\partial/\partial p_\mu\Big)^2$ to the sunset integral $\mathbf{J}(1,1,1,m)$, where the replacement $k_2 \to k_2+p$ is performed in (7). Further use is made of the relation

$$\Big(\partial/\partial p_\mu\Big)^2 P^{(i)}(p,m) = 4i(i+1-N/2)P^{(i+1)}(p,m) - 4i(i+1)m^2 P^{(i+2)}(p,m),$$

and the dimension property of the integral (see e.g. in Ref.[39]):

$$\left(p^2\frac{\partial}{\partial p^2} + m^2\frac{\partial}{\partial m^2}\right)\mathbf{J}(i,j,k,m) = (N-i-j-k)\,\mathbf{J}(i,j,k,m).$$

is taken into account. The third recurrence relation, which we need, has the following form:

$$\mathbf{J}(2,1,1,m) = \mathbf{J}(1,1,2,m) - \frac{2m^2}{\varepsilon}\mathbf{J}(1,1,3,m). \tag{15}$$

Eq.(15) is obtained in a similar manner as Eq.(14): the operator $\Big(\partial/\partial p_\mu\Big)^2$ is applied to the sunset $\mathbf{J}(1,1,1,m)$ integral with the initial distribution of momenta (7), and to the same diagram with the replacement $k_2 \to k_2+p$. Their difference yields the recurrence relation (15). We would like to note that Eq.(15) shows that the sunset diagram $\mathbf{J}(1,1,3,m)$ should be known up to terms of order ε^2.

After tedious calculation using representation (9), we obtain

$$\begin{aligned}
\mathbf{J}(1,2,2,m) &= \frac{K^2}{p^2}\Bigg(\log^2 y + 2\varepsilon\bigg[\frac{1}{2}\log^3 y - \zeta(2)\log y + \log(1-y)\log^2 y \\
&+2\log y\Big(3\mathrm{Li}_2(-y) + 2\mathrm{Li}_2(y)\Big) - 3\zeta(3) - 12\mathrm{Li}_3(-y) - 6\mathrm{Li}_3(y)\bigg]\Bigg) + O(\varepsilon^2), \\
\mathbf{J}(1,1,3,m) &= \frac{K^2}{m^2}\frac{1}{2(1-2\varepsilon)}\Bigg(\frac{1}{\varepsilon} + r\bigg[\log y + \varepsilon\bigg\{2\log y\log\left(\frac{y}{(1+y)^3(1-y)}\right) \\
&-\zeta(2) - 6\mathrm{Li}_2(-y) - 2\mathrm{Li}_2(y)\bigg\} + \varepsilon^2\bigg\{6\mathrm{S}_{1,2}(y^2) - 8\mathrm{S}_{1,2}(y) + 24\mathrm{S}_{1,2}(-y) \\
&-8\mathrm{Li}_3(y) - 24\mathrm{Li}_3(-y) - 11\zeta(3) - 2\zeta(2)\log\left(\frac{y^2}{(1+y)^3(1-y)}\right) \\
&+4\Big[\mathrm{Li}_2(y) + 3\mathrm{Li}_2(-y)\Big]\log\Big((1+y)^3(1-y)\Big) + 2\log y\log(1-y)\log\Big(y(1-y)\Big) \\
&+\frac{1}{6}\log^3 y + \frac{3}{2}\log y\log\left(\frac{y}{(1+y)^2}\right)\log\left(\frac{y}{(1+y)^6(1-y)^4}\right)\bigg\}\bigg] \\
&-\varepsilon\frac{3a+4}{2a}\bigg[\log^2 y - \varepsilon\bigg\{12\mathrm{Li}_3(y) + 24\mathrm{Li}_3(-y) + 6\zeta(3) \\
&+2\log y\Big(\zeta(2) - 4\mathrm{Li}_2(y) - 6\mathrm{Li}_2(-y)\Big) - 2\log^2 y\log(1-y) - \log^3 y\bigg\}\bigg]\Bigg) + O(\varepsilon^3),
\end{aligned}$$

where $r^2 = (4+a)/a$ and $y = (r-1)/(r+1)$. Putting $a=-1$, we get

$$\mathbf{J011}(1,2,2,m) = \frac{2}{3m^2}\biggl(\zeta(2) - \varepsilon\zeta(3) + \mathcal{O}(\varepsilon^2)\biggr), \tag{16}$$

and

$$\mathbf{J011}(1,1,3,m) = \frac{1}{2(1-2\varepsilon)m^2}\biggl(\frac{1}{\varepsilon} - \frac{\pi}{\sqrt{3}} + \varepsilon\biggl\{3\frac{\pi}{\sqrt{3}}\ln 3 - \frac{\zeta(2)}{3} - \frac{27}{2}S_2\biggr\}$$
$$+\varepsilon^2\biggl\{\frac{\zeta(3)}{3} - 27\frac{\mathrm{Ls}_3\left(\frac{2\pi}{3}\right)}{\sqrt{3}} - 14\frac{\pi}{\sqrt{3}}\zeta(2) - \frac{9}{2}\frac{\pi}{\sqrt{3}}\ln^2 3 + \frac{81}{2}S_2\ln 3\biggr\} + \mathcal{O}(\varepsilon^3)\biggr). \tag{17}$$

Note that the integrals **J011**(1,1,3,m) and **J011**(1,2,2,m) are much simpler than the master integrals (4) and (5).

3. *The* **F00111** *diagram.*

The diagram **F00111** demands special consideration. The investigations are similar to those done in [24]. Applying the integration-by-parts relation three times to one of the triangles in $\mathbf{F}(1,1,1,1,1,m)$ (every time with different distinctive line) and using the identity

$$2\mathbf{F}(1,1,2,1,1,m) + \mathbf{F}(1,1,1,1,2,m) = -\frac{d}{dm^2}\mathbf{F}(1,1,1,1,1,m),$$

we obtain the differential equation:

$$\biggl[(N-4)(p^4 + 2p^2m^2 + 3m^4) - 2m^2(p^4 + p^2m^2 + m^4)\frac{d}{dm^2}\biggr]\mathbf{F}(1,1,1,1,1,m) = f, \tag{18}$$

where the inhomogeneous term f contains only diagrams of $\mathbf{V_i}$, $\mathbf{J}$ and $\mathbf{L}$-type:

$$f = (p^4 - m^4)\biggl[\Bigl(\mathbf{L}(1,2,m) + \mathbf{L}(2,1,m)\Bigr)\mathbf{L}(1,1,m)$$
$$-\mathbf{V_1}(2,1,1,1,m) - \mathbf{V_2}(2,1,1,1,m)\biggr]$$
$$+3m^2(p^2+m^2)\mathbf{V_2}(1,2,1,1,m) - m^2(p^2-m^2)\mathbf{V_1}(1,2,1,1,m). \tag{19}$$

Expressing all **L**'s in terms of **L**(1,2,m) (see (8)) and using the recurrence relations for **V**-type integrals obtained via integration by parts, we derive a new, simpler representation for the inhomogeneous term f:

$$f \;=\; \frac{(p^2-m^2)}{(1-2\varepsilon)}\biggl[\frac{1-4\varepsilon}{1-2\varepsilon}\mathbf{T}_1^2(2,m) - \frac{p^2(3-8\varepsilon) + m^2(1-4\varepsilon)}{(1-2\varepsilon)}\mathbf{T}_1(2,m)\mathbf{L}(1,2,m)$$
$$+2(p^2+m^2)\mathbf{J}(1,2,2,m) + 4m^2\mathbf{J}(1,1,3,m) + 2m^2\mathbf{V_1}(1,1,2,1,m)\biggr]$$
$$+6m^2\Bigl[1 + (p^2-m^2)\frac{d}{dm^2}\Bigr]m^2\mathbf{V_2}(1,1,2,1,m), \tag{20}$$

where for the **V**-type diagrams the following differential equations hold:

$$\Big[(N-4)(p^2+m^2)-m^2-m^2(p^2+m^2)\frac{d}{dm^2}\Big]\mathbf{V_1}(1,1,2,1,m)$$
$$= m^2\mathbf{J}(1,2,2,m)+2(p^2-m^2)\Big[\mathbf{T}_1(3,m)\mathbf{L}(1,1,m)-\mathbf{J}(1,1,3,m)\Big], \qquad (21)$$
$$\Big[(N-6)p^2+3(N-4)m^2-2m^2(p^2+m^2)\frac{d}{dm^2}\Big]\mathbf{V_2}(1,1,2,1,m)$$
$$= -p^2\mathbf{J}(1,2,2,m)+3m^2\mathbf{T}_2(1,2,2,m), \qquad (22)$$

and (see e.g. Eq.(5.5) in Ref.[19])

$$\mathbf{T}_2(1,2,2,m)=\frac{K^2}{m^2}\cdot 3S_2+O(\varepsilon).$$

We integrate Eqs.(18), (21), (22) with the boundary condition that all diagrams tend to zero as $m^2\to\infty$. Finally for the on-shell case we obtain

$$\mathbf{F00111}(1,1,1,1,1,m)=8\int_0^1\frac{dx}{\sqrt{x(4-x)}}\arctan\sqrt{\frac{x}{4-x}}$$
$$\times\left\{\frac{\sqrt{3}}{1-x}\left(\frac{\pi}{6}-\arctan\frac{2x-1}{\sqrt{3}}\right)-\ln\left(1-x+x^2\right)\right\}.$$

This integral is evaluated numerically with a precision of 40 decimals. Then the program **PSLQ** (see above) is applied with 15 basis elements occurring in (1) to (6) with the result given in Table I.

Acknowledgments We are grateful to A. Davydychev, D. Kreimer and O. Veretin for useful discussions and careful reading of the manuscript and to O. Veretin for his help in numerical calculation. Two of us (J.F. and M.K.) are very indebted to O. V. Tarasov for useful comments. M.K. and A.K.'s research has been supported by the DFG project FL241/4-1 and in part by RFBR #98-02-16923.

References

[1] G. Degrassi, P. Gambino and A. Vicini, Phys. Lett. B **383** (1996) 219;
G. Degrassi, P. Gambino and A. Sirlin, Phys. Lett. B **394** (1997) 188.

[2] S. Bauberger and G. Weiglein, Phys. Lett. B **419** (1998) 333.

[3] P. Gambino, A. Sirlin and G. Weiglein, JHEP **9904** (1999) 025.

[4] F. Jegerlehner, Z. Phys. C **32** (1986) 425;
M. Consoli, W. Hollik and F. Jegerlehner, Phys. Lett. B **227** (1989) 167;
G. Degrassi, S. Fanchiotti and A. Sirlin, Nucl. Phys. B **351** (1991) 49;
G. Degrassi and A. Sirlin, Nucl. Phys. B **352** (1991) 342;
F. Halzen and B. A. Kniehl, Nucl. Phys. B **353** (1991) 567;
G. Degrassi, P. Gambino, M. Passera and A. Sirlin, Phys. Lett. B **418** (1997) 209.

[5] V. A. Smirnov, Phys. Lett. B **394** (1997) 205;
A. Czarnecki and V. A. Smirnov, Phys. Lett. B **394** (1997) 211;
L .V. Avdeev and M. Yu. Kalmykov, Nucl. Phys. B **502** (1997) 419.

[6] O V. Tarasov, Nucl. Phys. B **502** (1997) 455.

[7] J. Fleischer and M. Yu. Kalmykov, in preparation.

[8] A. A. Vladimirov, Theor. Math. Phys. **43** (1980) 417.

[9] F. V. Tkachov, Phys. Lett. B **100** (1981) 65;
K. G. Chetyrkin and F. V. Tkachov, Nucl. Phys. B **192** (1981) 159.

[10] A. N. Vassiliev, Y. M. Pis'mak and Y. P. Honkonen, Theor. Math. Phys. **47** (1981) 465.

[11] J. van der Bij and M. Veltman, Nucl. Phys. B **231** (1984) 205;
F. Hoogeveen, Nucl. Phys. B **259** (1985) 19;
J. van der Bij and F. Hoogeveen, Nucl. Phys. B **283** (1987) 477.

[12] N. Gray, D. J. Broadhurst, W. Grafe and K. Schilcher, Z. Phys. C **48** (1990) 673;
D. J. Broadhurst, N. Gray and K. Schilcher, Z. Phys. C **52** (1991) 111;
D. J. Broadhurst, Z. Phys. C **54** (1992) 599.

[13] A. V. Kotikov, Phys. Lett. B **254** (1991) 158; B **259** (1991) 314; Phys. Lett. B **267** (1991) 123; B **295** (1992) 409(E); JHEP **9809** (1998) 001.

[14] A. V. Kotikov, Mod. Phys. Lett. A **6** (1991) 677.

[15] C. Ford, I. Jack and D. R. T. Jones, Nucl. Phys. B **387** (1992) 373; B **504** (1997) 551(E).

[16] U. Nierste, D. Müller and M. Böhm, Z. Phys. C **57** (1993) 605;
A. I. Davydychev and N. I. Ussyukina, Phys. Lett. B **332** (1994) 159;
F. A. Berends, A. I. Davydychev and V. A. Smirnov, Nucl. Phys. B **478** (1996) 59;
A. I. Davydychev and R. Delbourgo, J. Math. Phys. **39** (1998) 4299;
J. Fleischer, F. Jegerlehner, O. V. Tarasov and O. L. Veretin, Nucl. Phys. B **539** (1999) 671.

[17] D. J. Broadhurst, Z. Phys.C **47** (1990) 115.

[18] D. J. Broadhurst, J. Fleischer and O. V. Tarasov, Z. Phys. C **60** (1993) 287.

[19] A. I. Davydychev and J. B. Tausk, Nucl. Phys. B **397** (1993) 123.

[20] A. Djouadi, Nuovo Cim. A **100** (1988) 357;
P. N. Maher, L. Durand and K.Riesselmann, Phys. Rev. D **48** (1993) 1061; D **52** (1995) 553(E);
R. Scharf and J. B. Tausk, Nucl. Phys. B **412** (1994) 523.

[21] F. A. Berends, M. Buza, M. Böhm and R. Scharf, Z. Phys.C **63** (1994) 227;

F. A. Berends and J. B. Tausk, Nucl. Phys. B **421** (1994) 456;

S. Bauberger, F. A. Berends, M. Böhm and M. Buza, Nucl. Phys. B **434** (1995) 383.

[22] A. Ghinculov and J. J. van der Bij, Nucl. Phys. B **436** (1995) 30;

F. A. Berends, A. I. Davydychev, V. A. Smirnov and J. B. Tausk, Nucl. Phys. B **439** (1995) 536;

A. Ghinculov and Y. -P. Yao, Nucl. Phys. B **516** (1998) 385;

A. T. Suzuki and A. G. M. Schmidt, Eur. Phys. J. C **5** (1998) 175.

[23] V. Borodulin and G. Jikia, Phys. Lett. B **391** (1997) 434.

[24] J. Fleischer, A. V. Kotikov and O. L. Veretin, Nucl. Phys. B **547** (1999) 343.

[25] A. I. Davydychev and J. B. Tausk, Phys. Rev. D **53** (1996) 7381.

[26] F. A. Berends, A. I. Davydychev and N. I. Ussyukina, Phys. Lett. B **426** (1998) 95.

[27] A. I. Davydychev and A. G. Grozin, Phys. Rev. D **59** (1999) 054023.

[28] D. Kreimer, Phys. Lett. B **273** (1991), 277; ibid B **292** (1992), 341;

A. Czarnecki, U. Kilian and D. Kreimer, Nucl. Phys. B **433** 1995) 259;

S. Bauberger, and M. Böhm, Nucl. Phys. B **445** (1995) 25.

[29] S. Groote, J. G. Körner and A. A. Pivovarov, Phys. Lett. B **443** (1998) 269; Nucl. Phys. B **542** (1999) 515.

[30] D. J. Broadhurst, Z. Phys. C **54** (1992) 599.

[31] L. Lewin, *Polylogarithms and associated functions* (North-Holland, Amsterdam, 1981).

[32] J. Fleischer and O. V. Tarasov, Z. Phys. C **64** (1994) 413;

O. V. Tarasov, Nucl. Phys. B **480** (1996) 397.

[33] L. V. Avdeev, J. Fleischer, M. Yu. Kalmykov and M. N. Tentyukov, Nucl. Inst. Meth. A **389** (1997) 343; Comp. Phys. Commun. **107** (1997) 155.

[34] J. Fleischer, Int. J. Mod. Phys. C **6** (1995) 495;

J. Fleischer, V. A. Smirnov and O. V. Tarasov, Z. Phys. C **74** (1997) 379;

J. Fleischer et al., Eur. Phys. J. C **2** (1998) 747;

J. Fleischer, M. Yu. Kalmykov and O. L. Veretin, Phys. Lett. B **427** (1998) 141.

[35] D. J. Broadhurst, Eur. Phys. J. C **8** (1999) 311.

[36] H. R. P. Ferguson, D. H. Bailey and S. Arno, NASA-Ames Technical Report, NAS-96-005;

D. H. Bailey and D. J. Broadhurst, math.NA/9905048.

[37] J. Fleischer, A. V. Kotikov and O. L. Veretin, Phys. Lett. B **417** (1998) 163.

[38] A. I. Davydychev and V. A. Smirnov, hep-ph/9903328, to appear in Nucl. Phys. B.

[39] M. Gaffo, H. Czyz, S. Laporta and E. Remiddi, Nuovo Cim. A **111** (1998) 365.



Figure 1: The full set of two-loop self-energies diagrams with one mass. Bold and thin lines correspond to the mass and massless propagators, respectively.